

Elementary divisors of Gram matrices of certain Specht modules

(jt. w. G. Nebe and A. Mathas)

Notation

\mathcal{S}_n	symmetric group
$\lambda \vdash n$	partition of n
λ'	transpose of λ
S^λ	integral Specht module over $\mathbf{Z}\mathcal{S}_n$
$n_\lambda := \text{rk}_{\mathbf{Z}} S^\lambda$	
$S^{\lambda,*}$	\mathbf{Z} -linear dual
$S_A^\lambda := A \otimes_{\mathbf{Z}} S^\lambda$	for a commutative ring A
$(=, -)$	the \mathcal{S}_n -invariant nondegenerate \mathbf{Z} -bilinear form on S^λ

Problem

$\mathbf{Z}\mathcal{S}_n$ -linear map:

$$\begin{array}{ccc}
 S^\lambda & \xhookrightarrow{\eta} & S^{\lambda,*} \\
 \xi & \longmapsto & (\xi, -)
 \end{array}$$

Determine quotient $S^{\lambda,*}/S^\lambda$ as a finite abelian group.

Equivalently, determine el. div. of Gram matrix of $(=, -)$.

Motivation

I) Simple modules

Ordinary:

$$\{S_{\mathbf{C}}^{\lambda} \mid \lambda \vdash n\} = \{\text{simple } \mathbf{C}\mathcal{S}_n\text{-modules}\}$$

Modular: p prime.

$$D_{\mathbf{F}_p}^{\lambda} := \text{Im} \left((S^{\lambda} \xrightarrow{\eta} S^{\lambda,*}) \otimes_{\mathbf{Z}} \mathbf{F}_p \right)$$

James proves:

$$\{D_{\mathbf{F}_p}^{\lambda} \mid \lambda \vdash n \text{ } p\text{-regular}\} = \{\text{simple } \mathbf{F}_p\mathcal{S}_n\text{-modules}\}$$

So

$$\dim_{\mathbf{F}_p} D_{\mathbf{F}_p}^{\lambda} = \#\{\text{el. div. of } S^{\lambda} \text{ not divisible by } p\}$$

However, we will make extensive use of known such dimensions rather than derive them.

II) Quasiblocks

How to understand the integral group ring $\mathbf{Z}\mathcal{S}_n$?

Wedderburn isomorphism:

$$\mathbf{Q}\mathcal{S}_n \xrightarrow{\sim} \prod_{\lambda \vdash n} \underbrace{\text{End}_{\mathbf{Q}} S_{\mathbf{Q}}^\lambda}_{\simeq \mathbf{Q}^{n_\lambda \times n_\lambda}}$$

Restrict to Wedderburn embedding.

$$\mathbf{Z}\mathcal{S}_n \hookrightarrow \prod_{\lambda \vdash n} \underbrace{\text{End}_{\mathbf{Z}} S^\lambda}_{\simeq \mathbf{Z}^{n_\lambda \times n_\lambda}}$$

Describe $\mathbf{Z}\mathcal{S}_n$ as image of the Wedderburn embedding.

Example:

$$\mathbf{Z}\mathcal{S}_3 \simeq \mathbf{Z} \begin{array}{c} \xrightarrow{3} \\ \xrightarrow{3} \\ \xrightarrow{2} \end{array} \left(\begin{array}{cc} \mathbf{Z} & \mathbf{Z} \\ 3\mathbf{Z} & \mathbf{Z} \end{array} \right) \begin{array}{c} \xrightarrow{3} \\ \xrightarrow{3} \\ \xrightarrow{2} \end{array} \mathbf{Z}$$

Call $Q^{(2,1)} = \begin{pmatrix} \mathbf{Z} & \mathbf{Z} \\ 3\mathbf{Z} & \mathbf{Z} \end{pmatrix}$ a *quasiblock* of $\mathbf{Z}\mathcal{S}_3$.

Orthogonal decomposition into rational primitive central idempotents:

$$1_{\mathbf{Q}\mathcal{S}_n} = \sum_{\lambda \vdash n} \varepsilon_\lambda$$

General definition of a *quasiblock* :

$$Q^\lambda := \mathbf{Z}\mathcal{S}_n \varepsilon_\lambda \left(\hookrightarrow \text{End}_{\mathbf{Z}} S^\lambda \simeq \mathbf{Z}^{n_\lambda \times n_\lambda} \right)$$

Unknown: index of this inclusion (as abelian groups).

Necessary condition on $\varphi \in \text{End}_{\mathbf{Z}} S^\lambda$ to lie in image:

$$\eta^{-1} \varphi \eta \quad \text{has to be integral}$$

(η^{-1} = rational inverse of η). Not sufficient in general.

The Smith form of $S^{(2,1)}$ is $\begin{pmatrix} 1 & 0 \\ 0 & 3 \end{pmatrix}$, so for $Q^{(2,1)} = \begin{pmatrix} \mathbf{Z} & \mathbf{Z} \\ 3\mathbf{Z} & \mathbf{Z} \end{pmatrix}$ this necessary condition is in fact sufficient.

In general, elementary divisors of S^λ partially describe Q^λ .

Method

I) Jantzen filtration

Let p be a prime, let $i \geq 0$. Define $S_{\mathbf{F}_p}^\lambda(i)$ by

$$\begin{array}{ccccc} \text{Kern} & \hookrightarrow & S^\lambda & \xrightarrow{\eta} & S^{\lambda,*}/p^i S^{\lambda,*} \\ \downarrow & & \downarrow & & \\ S_{\mathbf{F}_p}^\lambda(i) & \hookrightarrow & S_{\mathbf{F}_p}^\lambda & & \end{array}$$

The *Jantzen filtration* is the finite filtration

$$S_{\mathbf{F}_p}^\lambda = S_{\mathbf{F}_p}^\lambda(0) \supseteq S_{\mathbf{F}_p}^\lambda(1) \supseteq S_{\mathbf{F}_p}^\lambda(2) \supseteq \cdots,$$

We have $S_{\mathbf{F}_p}^\lambda(0)/S_{\mathbf{F}_p}^\lambda(1) \simeq D_{\mathbf{F}_p}^\lambda$.

The Jantzen filtration gives $\mathbf{Z}_{(p)}$ -linear elementary divisors:

$$S_{\mathbf{Z}_{(p)}}^{\lambda,*}/S_{\mathbf{Z}_{(p)}}^\lambda \simeq \bigoplus_{i \geq 0} (\mathbf{Z}/p^i \mathbf{Z})^{\dim_{\mathbf{F}_p} S_{\mathbf{F}_p}^\lambda(i)/S_{\mathbf{F}_p}^\lambda(i+1)}.$$

So to get elementary divisors:

- Determine composition factors of Jantzen subquotients (using Jantzen's Lemma and Schaper's Theorem).
- Use results of James et al. on the dimensions of simple modules.

II) Jantzen's Lemma

p prime, $\lambda \vdash n$ arbitrary, $\mu \vdash n$ p -regular.

$$\vartheta_i := [S_{\mathbf{F}_p}^\lambda(i)/S_{\mathbf{F}_p}^\lambda(i+1) : D_{\mathbf{F}_p}^\mu].$$

Jantzen's Lemma:

$$\begin{aligned} [S_{\mathbf{F}_p}^\lambda : D_{\mathbf{F}_p}^\mu] &= \vartheta_0 + \vartheta_1 + \vartheta_2 + \cdots \\ [S_{\mathbf{Z}_{(p)}}^{\lambda,*}/S_{\mathbf{Z}_{(p)}}^\lambda : D_{\mathbf{F}_p}^\mu] &= 0\vartheta_0 + 1\vartheta_1 + 2\vartheta_2 + \cdots \end{aligned}$$

So if $[S_{\mathbf{F}_p}^\lambda : D_{\mathbf{F}_p}^\mu] = 1$, then

$$\vartheta_i = \begin{cases} 1 & \text{for } i = [S_{\mathbf{Z}_{(p)}}^{\lambda,*}/S_{\mathbf{Z}_{(p)}}^\lambda : D_{\mathbf{F}_p}^\mu] \\ 0 & \text{else.} \end{cases}$$

If $\lambda = \mu$, then $\vartheta_0 = 1$, and $\vartheta_i = 0$ for $i > 0$.

If $\lambda \neq \mu$, we have $\vartheta_0 = 0$. E.g. if $[S_{\mathbf{F}_p}^\lambda : D_{\mathbf{F}_p}^\mu] = 2$, then:

If $[S_{\mathbf{Z}_{(p)}}^{\lambda,*}/S_{\mathbf{Z}_{(p)}}^\lambda : D_{\mathbf{F}_p}^\mu] = 2$, then $\vartheta_1 = 2$.

If $[S_{\mathbf{Z}_{(p)}}^{\lambda,*}/S_{\mathbf{Z}_{(p)}}^\lambda : D_{\mathbf{F}_p}^\mu] = 3$, then $\vartheta_1 = 1$, $\vartheta_2 = 1$.

If $[S_{\mathbf{Z}_{(p)}}^{\lambda,*}/S_{\mathbf{Z}_{(p)}}^\lambda : D_{\mathbf{F}_p}^\mu] = 4$, then $\vartheta_1 = 1$, $\vartheta_3 = 1$; or $\vartheta_2 = 2$.

Etc. So there is still ambiguity.

III) Schaper's Theorem

Even if all occurring decomposition numbers $[S_{\mathbf{F}_p}^\lambda : D_{\mathbf{F}_p}^\mu]$ are known, we still need $[S_{\mathbf{Z}(p)}^{\lambda,*} / S_{\mathbf{Z}(p)}^\lambda : D_{\mathbf{F}_p}^\mu]$.

Schaper's Theorem is a formula in the Grothendieck group:

$$[S_{\mathbf{Z}(p)}^{\lambda,*} / S_{\mathbf{Z}(p)}^\lambda] = \sum_{\lambda \trianglelefteq \nu} \alpha_\nu [S_{\mathbf{F}_p}^\nu],$$

with combinatorially determined coefficients $\alpha_\nu \in \mathbf{Z}$.

So for our purposes, knowledge of further decomposition numbers $[S_{\mathbf{F}_p}^\nu : D_{\mathbf{F}_p}^\mu]$ is required.

IV) Transposition

Jantzen filtration of $S^\lambda \longleftrightarrow$ Jantzen filtration of $S^{\lambda'}$

Some results

Table of calculated Jantzen filtrations.

Partially depending on conjectures on dec. numbers.

Partially with ambiguity.

λ	Remarks
$(n - m, 1^{n-m})$	hooks, two methods: Jantzen-Schaper and direct; independently done by James-Mathas
$(n - m, m)$	two-row; independently done by Fayers and by Murphy
$(n - 3, 2, 1)$	
$(n - 4, 2^2)$	if $p = 2$: depending on a conj. on $[S_{\mathbf{F}_2}^{(n-4,2^2)} : D_{\mathbf{F}_2}^{(n)}]$; if $p = 3$: with ambiguity
$(n - 4, 3, 1)$	if $p = 2$: with ambiguity
$(n - 4, 2, 1^2)$	if $p = 2$: depending heavily on conj. on dec. nos., and with ambiguity; if $p = 3$: depending on conj. on dec. nos.
Case $n - \lambda_1 < p$	using Kleshchev's Modular Branching and Carter-Payne

How to proceed?

Let e^μ be a primitive idempotent of $\mathbf{Z}_{(p)}\mathcal{S}_n$ belonging to $D_{\mathbf{F}_p}^\mu$ and let ε^λ be the primitive central idempotent of $\mathbf{Q}\mathcal{S}_n$ belonging to $S_{\mathbf{Q}}^\lambda$.

The embedding η decomposes into a direct sum of embeddings

$$S_{\mathbf{Z}_{(p)}}^\lambda e^\mu \hookrightarrow^{e^\mu \eta e^\mu} S_{\mathbf{Z}_{(p)}}^{\lambda,*} e^\mu ,$$

so we get a block diagonalisation of the Gram matrix, with blocks of edge length

$$\text{rk}_{\mathbf{Z}_{(p)}} S^\lambda e^\mu = [S_{\mathbf{F}_p}^\lambda : D_{\mathbf{F}_p}^\mu] .$$

It remains to

calculate the Smith form of $e^\mu \eta e^\mu$.

Schaper's Theorem ess. allows to calculate the determinant:

$$v_p(\det(e^\mu \eta e^\mu)) = [S_{\mathbf{Z}_{(p)}}^{\lambda,*} / S_{\mathbf{Z}_{(p)}}^\lambda : D_{\mathbf{F}_p}^\mu]$$

Assume $[S_{\mathbf{F}_p}^\lambda : D_{\mathbf{F}_p}^\mu] \geq 1$. Let

$$Q_{(p)}^{\lambda:\mu} := e^\mu Q_{\mathbf{Z}_{(p)}}^\lambda e^\mu \left(\hookrightarrow \mathbf{Z}_{(p)}^{[S_{\mathbf{F}_p}^\lambda : D_{\mathbf{F}_p}^\mu] \times [S_{\mathbf{F}_p}^\lambda : D_{\mathbf{F}_p}^\mu]} \right).$$

- $\mathbf{Q} \otimes_{\mathbf{Z}_{(p)}} Q_{(p)}^{\lambda:\mu}$ is simple.
- $Q_{(p)}^{\lambda:\mu}$ is local (for $e^\mu Q_{\mathbf{Z}_{(p)}}^\lambda$ is indecomposable).
- $e^\mu \eta e^\mu$ is an inclusion of simple $Q_{(p)}^{\lambda:\mu}$ -lattices.

If we could classify all inclusions of simple lattices over $Q_{(p)}^{\lambda:\mu}$, we would know the Smith form of $e^\mu \eta e^\mu$ to be one of the Smith forms of these inclusions (moreover, one of a given determinant).

Example: $Q_{(2)}^{(3,1^2):(5)} \simeq \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathbf{Z}_{(2)}^{2 \times 2} \mid b \equiv_2 0, a \equiv_2 d \right\}$,
with simple lattices $X := \begin{pmatrix} \mathbf{Z}_{(2)} \\ \mathbf{Z}_{(2)} \end{pmatrix}$ and $Y := \begin{pmatrix} 2\mathbf{Z}_{(2)} \\ \mathbf{Z}_{(2)} \end{pmatrix}$.

Inclusions of determinant 4: only the scalar inclusions $X \xrightarrow{2} X$ and $Y \xrightarrow{2} Y$, both with Smith form $\begin{pmatrix} 2 & 0 \\ 0 & 2 \end{pmatrix}$.

Need: lattices over $Q_{(p)}^{\lambda:\mu}$. Note: $\text{rk}_{\mathbf{Z}_{(p)}} Q_{(p)}^{\lambda:\mu} = [S_{\mathbf{F}_p}^\lambda : D_{\mathbf{F}_p}^\mu]^2$.

So to ask for $Q_{(p)}^{\lambda:\mu}$ can be viewed a refined version of the question for the decomposition numbers.

Elementary divisors over $\mathbf{Z}[q, q^{-1}]$?

For the Specht module over the Hecke algebra \mathcal{H} with ground ring $\mathbf{Z}[q, q^{-1}]$, we may also consider the Gram matrix of the invariant bilinear form.

Distinguish (also over $\mathbf{Z}_{(p)}[q, q^{-1}]$):

- divisibly diagonalizable matrices (each diagonal entry divides successor)
- diagonalizable, but not divisibly diagonalizable matrices
- non-diagonalizable matrices

Andersen remarked that not every Gram matrix of a Specht module is diagonalizable.

For $\lambda = (n - l, 1^l)$: bases divisibly diagonalizing the Gram matrix (the q causing difficulties).

For $\lambda = (3, 3, 2)$: not diagonalizable over $\mathbf{Z}_{(2)}[q, q^{-1}]$.

For $\lambda = (4, 2, 1, 1)$: neither diagonalizable over $\mathbf{Z}_{(2)}[q, q^{-1}]$ nor over $\mathbf{Z}_{(3)}[q, q^{-1}]$.

Method: if diagonalizable, then there is a connection between the elementary divisors over $\mathbf{Q}[q, q^{-1}]$ and $\mathbf{F}_p[q, q^{-1}]$.

We do not know (not even conjecturally) a criterion for λ to decide on diagonalizability.